



A multi-faceted enzyme lab:

Looking at the effect of an organophosphate insecticide on acetylcholinesterase activity in the bean beetle

Hector Fermin, Gurcharan Singh, Rahat Gul, Min Gyu Noh, Violeta Contreras Ramirez, and Fardad Firooznia

City College of New York, New York, NY 10031

Why use this system?

- Bean beetles are common pests
- Bean beetles are easy to maintain in the laboratory
- Organophosphate insecticides are commonly used to control this pest organism
- Ties in several course topics: Introduction to enzymes, the role of inhibitors in enzyme activity, cell-cell signaling, and potential ecological consequences



Materials

- Bean beetles, *Callosobruchus maculatus*
- Incubator with light and temperature control
- Insecticide malaoxon
- ATCI as the substrate
- DTNB as the indicator
- Spectrophotometer, vortexer, centrifuge
- Water bath, autoclave, micropipettors
- Other standard laboratory equipment



What should students know?

- An introduction to the life cycle of the insects
- Basic structure of proteins
- Basics about enzymes
- The basics of competitive and non-competitive inhibition
- Basics of synaptic signaling
- Function of acetylcholinesterase




Image source:

http://www.cnsforum.com/imagebank/item/rcpt_sys_ACh_esterase/default.aspx
used with permission


Sample Student Results



Table 1: Relative inhibition of activity of AChE (acetylcholinesterase) caused by the insecticide malaoxon. AChE was extracted from bean beetles grown on black-eyed peas. Relative inhibition calculated as:
 $100 \times \frac{(\text{Absorbance without malaoxon} - \text{Absorbance with malaoxon})}{\text{Absorbance without malaoxon}}$


| Beetle # | Gender | Relative inhibition (%) |
|----------|--------|-------------------------|
| 1 | F | 49.6 |
| 2 | M | 30.2 |
| 3 | M | 34.2 |
| 4 | M | 17.7 |
| 5 | M | 25 |
| 6 | F | 41.2 |
| 7 | F | 12.5 |
| 8 | M | 3.42 |
| Mean | | 26.7 |
| SD | | 15.3 |

Figure 2: The effect of different concentrations of the insecticide malaoxon on the AChE enzyme activity. Columns represent mean relative inhibition caused by the insecticide, and error bars represent standard deviations. Sample size was 4.




Enzyme Assay


Based on the work done by Ellman et al. (1961), Ffrench-Constant and Bonning (1989), Spencer et al. (1998), and Gbaye et al. (2012). In the presence of the enzyme AChE, the compound ATCI (acetylthiocholine iodide) is hydrolyzed to produce acetate and thiocholine.




The thiol (R-SH) group of thiocholine can react with the indicator compound DTNB (5,5'-dithio-bis-2-nitrobenzoic acid) to form TNB⁻ (5-thio-2-nitrobenzoate), which ionizes to the TNB²⁻ in water at mildly alkaline pH; this is yellow and can be detected spectrophotometrically.



Basic Procedure



Procedure for Comparing Different Concentrations of Malaoxon



Comments



- This lab exercise is easily doable with first semester college students
- It can be done in regular 3-hour laboratories
- The topic can be supplemented with readings such as:
 - Rachel Carson's "Silent Spring"
 - London et al. (2005) linking organophosphate insecticides and suicide
 - Sadeghi Hashjin et al. (2013) linking malathion with anxiety in rodents

Current Research

- To develop longer-term student projects to determine whether food source affects sensitivity of the AChE to insecticide
- To determine whether the beetles can be frozen at -20°C or -80°C for future studies without significant changes in enzyme activity
- To determine whether incubation time can be optimized

Acknowledgements

- Biology Department and Division of Science, CCNY
- Mariano Soley (lab instructor) and Students who participated in this project
- Chris Beck and Larry Blumer for beetles, instructions, beetle images, and other help
- Funding by NSF DUE-0815135, and DUE-0814373 to Morehouse College and Emory University
- **Disclaimer:** Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation

References

- Beck, C. W., and Blumer, L. S. (2011). A handbook on bean beetles, *Callosobruchus maculatus*. www.beanbeetles.org.
- Carson, R. (1962). Silent Spring. Mariner Books.
- Ellman, G. L., Courtney, K. D., Andres, V., and Featherstone, R. M. (1961). A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology*, 7: 88-95.
- Ffrench-Constant, R. H., and Bonning, B. C. (1989). Rapid microtitre plate test distinguishes insecticide resistant acetylcholinesterase genotypes in the mosquitoes *Anopheles albimanus*, *An. nigerrimus* and *Culex pipiens*. *Medical and Veterinary Entomology*, 3: 9-16.
- Gbaye, O. A., Holloway, G. J., and Callaghan, A. (2012). Variation in the sensitivity of *Callosobruchus* (Coleoptera: Bruchidae) acetylcholinesterase to the organophosphate insecticide malaoxon: effect of species, geographical strain and food type. *Pest Management Science*, 68: 1265-1271.
- Longon, L., Flisher, A. J., Wesseling, C., Mergler, D., and Kromhout, H. (2005). Suicide and Exposure to Organophosphate Insecticides: Cause or Effect? *American Journal of Industrial Medicine*, 47:308-321.
- Sadeghi Hashjin, G., Sadeghi Dizaj, F., Attaran, H., and Koochi, M. K. (2013). Malathion induces anxiety in the male adult mouse. *Archives of Medical Science*, 9(2): 368-371.
- Spencer, A. G., Price, N. R., and Callaghan, A. (1998). Malathion-specific resistance in a strain of the rust red grain beetle *Cryptolestes ferrugineus* (Coleoptera: Cucujidae). *Bulletin of Entomological Research*, 88: 199-206.